Barb

Barbara K. Zedler, M.D.
Director, Clinical Operations
Clinical Evaluation Group
Worldwide Scientific Affairs
Philip Morris USA
Richmond, VA
barbara.k.zedler@pmusa.com
804.274.2799
cell 804.651.3544

-----Original Message-----
From: Roethig, Hans
Sent: Monday, October 13, 2003 4:42 PM
To: Nelson, Bettie L; Zedler, Barbara; Kinser, Robin D.; Feng, Shixia
Cc: Feng, Shixia
Subject: RE: CREATININE CLEARANCE ADJUSTMENT

But we do not calculate the creatinine clearance.

-----Original Message-----
From: Nelson, Bettie L
Sent: Monday, October 13, 2003 11:50 AM
To: Zedler, Barbara; Kinser, Robin D.; Feng, Shixia
Cc: Feng, Shixia; Roethig, Hans
Subject: CREATININE CLEARANCE ADJUSTMENT

Below is the Cockcroft and Gualt formula that MDS uses for creatinine clearance to control for gender specific creatinine.
The calculated creatinine clearance will be determined by the laboratory, based on the serum creatinine, age, and body weight in kg, using the Cockcroft-Gault formula.

$$\frac{(140 - \text{age [years]}) \times \text{body weight [kg]} \times K}{\text{serum creatinine [mol/L]}}$$

$$1.73\ m^2$$

K = 1.23 for men, 1.05 for women

Please let me know whether you have any changes to this method for adjusting creatinine.

With best regards,

Bettie

Source: https://www.industrydocuments.ucsf.edu/docs/tnpj0001

(804) 274-1502

(804) 334-5923 (Cellular)

(804) 274-3982 (Fax)

PM3001373820

Source: https://www.industrydocuments.ucsf.edu/docs/tnpj0001